arXiv:1207.0930v2 [math.AG] 6 Aug 2012

CONSTRUCTIVE DEGREE BOUNDS FOR GROUP–BASED MODELS

MATEUSZ MICHAŁEK

Abstract. Group–based models arise in algebraic statistics while studying evolution processes. They are represented by embedded algebraic varieties, that turn out to be toric. Both from the theoretical and applied point of view one is interested in determining the ideals defining the varieties. Conjectural bounds on the degree in which these ideals are generated were given by Sturmfels and Sullivant [SS05, Conjectures 29, 30]. We prove that for the 3–Kimura model, corresponding to the group $G = \mathbb{Z}_2 \times \mathbb{Z}_2$, the projective scheme can be defined by an ideal generated in degree 4. In particular, it is enough to consider degree 4 phylogenetic invariants to test if a given point belongs to the variety. Moreover, we give a constructive proof, that for any fixed, finite abelian group G there exists a constant d, such that for any tree the associated projective scheme can be defined by an ideal generated in degree at most d.

1. Introduction

We investigate properties of a special class of not necessarily normal, projective toric varieties. Their construction is motivated by applied mathematics, namely phylogenetics. We will not describe the connections with applications, referring to [SS03], [PS05]. On the other hand we give an exact mathematical definition of the class of varieties we consider. We present the problems in toric geometry, arising from applied mathematics, so that pure mathematicians can address them.

Let T be a rooted tree, that is a connected graph without cycles and with one distinguished vertex v_0 called the root. We direct the edges of T away from the root. Let V be the set of vertices of T. For each $v \in V$, consider a vector space W_v with a fixed basis $e^v_1, \dots, e^v_{n(v)}$. We will construct subvarieties in $\bigotimes_{v\in L} W_v$, where L is the set of leaves of T. The variety is given as a closure of the parametrization map

$$\phi : W_{v_0} \times \prod_{(v_1,v_2)} W^*_{v_1} \otimes W_{v_2} \to \bigotimes_{v\in L} W_v,$$

where the product $\prod_{(v_1,v_2)}$ is taken over all edges of T. Consider the basis $\bigotimes_{v\in V} e^v_{j(v)}$ of the vector space $\bigotimes_{v\in L} W_v$, indexed by functions j that associate to a vertex v an index from 1 to $n(v)$. We define ϕ by:

$$(\bigotimes_{v\in V} e_{j(v)})^*(\phi((w_0, \prod_{(v_1,v_2)} w_{v_1^*,v_2}))) = \sum_i (e^{v_0}_{i(v_0)})^*(w_0) \prod_{(v_1,v_2)} (e_{i(v_1)} \otimes e^*_{i(v_2)})(w_{v_1^*,v_2}),$$

where $w_{v_1^*,v_2} \in W^*_{v_1} \otimes W_{v_2}$ and the sum $\sum_i$ is taken over all functions that associate to a vertex v an index of a basis vector of W_v and agree with j, when v is a leaf. As usually by * we denote the dual vector, with respect to the chosen basis.

2010 *Mathematics Subject Classification.* 14M25, 52B20.

Key words and phrases. group-based model, 3-Kimura, phylogenetic invariant, toric variety.

The author is supported by the NCN grant UMO.

The image of ϕ is called *the general Markov model.* Let $K_{1,n}$ be the *claw tree* – cf. Definition 3.2, with the inner vertex v_0. We encourage the reader to check that for this tree and W_{v_0} one dimensional, the obtained variety is the Segre embedding of $V_1,\dots,V_n$. Moreover, when W_{v_0} is k dimensional, the Zariski closure of the image is the k-th secant variety of the Segre variety. More details on this construction, including motivation, can be found in [PS05, Chapter 4] or a short paper [ERSS04].

We are interested in equivariant versions of the above construction. First examples of such appeared in phylogenetics, the science that aims at reconstructing the evolution of DNA [Kim80], [Kim81], [JC69]. It was observed that certain symmetries between nucleobasis of DNA can be induced by an action of a group. Important mathematical implications of such a description of the variety were proven in [HP89][1]. Along with other works, [ES93, SSE93], they proved that, using modern language, if the group is sufficiently big, the obtained variety is toric – we do not assume that toric varieties have to be normal. Such models are called *group–based* and are the central object of study of this article.

In this setting we assume that all vector spaces $W_v \cong W$ are regular representations of a fixed finite abelian group G. Thus we can identify the basis with group elements. We restrict the domain of ϕ to G-invariant vectors. More precisely we consider:

$$\phi_G : W^G \times \prod_{(v_1,v_2)} (W^* \otimes W)^G \to \bigotimes_{v\in L} W,$$

where the action of the group on the tensor product and on the dual is induced from the vector space. Note that W^G is one dimensional, hence the image coincides with the image of $\prod_{(v_1,v_2)}(W^*\otimes W)^G$. The fact that W^G is one dimensional corresponds in biology to the assumption of *uniform root distribution.* The Zariski closure of the image will be denoted by $X(T,G)$ and is known to be toric. We also let $X_{\mathbb{P}}(T,G)$ be the projective variety, such that $X(T,G)$ is the affine cone over $X_{\mathbb{P}}(T,G)$.

More generally, one can consider the vector spaces W_v to be arbitrary representations. The model in such a case is called *equivariant* – it was introduced in [DK09]. This construction does not always lead to a toric variety. Indeed, the first presented example – secants of Segre – is an equivariant model with the trivial group.

The study of equivariant models was reduced to the case of claw trees. Still, the description of the varieties associated to claw trees is difficult. Let us present what is known.

1. The equivariant model associated to the trivial group corresponds to secant variety of the Segre embedding. These varieties are an object of intensive studies. The description of the first secant was recently given in [Rai11], solving the GSS conjecture. It is known that analogous description, by flattenings, for higher secants does not hold. On the other hand, it was proven that, for a fixed k, there exists d, such that the k-th secant of the Segre variety, for any number of factors, is set theoretically described by equations of degree at most d, [DK11].
2. The group–based model with $G = \mathbb{Z}_2$ is the only model where we can explicitly give a description of the ideal [SS05, Theorem 28], [CP07]. It is definitely the simplest model one can consider. However, it turned out to be very interesting due to nice properties [BW07] and connections with different branches of mathematics [SX10, Man09]. For the generalizations of the construction to graphs we advise the reader to consult [BBKM10].

---

[1] We would like to thank Elizabeth Allman for brigning this fact to our attention.

(3) Equivariant models for an abelian group G include previous examples. Recently, it was proved in [DE] that set theoretically they can be defined in (some) bounded degree.

The following conjecture is due to Sturmfels and Sullivant.

Conjecture 1.1 (Conjecture 30)**.** *The ideal for the 3-Kimura model, that is a group–based model for* $G = \mathbb{Z}_2 \times \mathbb{Z}_2$*, is generated in degree at most* 4 *for any tree* T*.*

One is particulary interested in this model, as it was introduced by theoretical biologists.

Main Theorem 1. *For any tree* T *the projective scheme* $X_{\mathbb{P}}(T, \mathbb{Z}_2 \times \mathbb{Z}_2)$ *can be defined by an ideal generated in degree at most* 4*.*

Conjecture 1.1 has the following generalization, also due to Sturmfels and Sullivant.

Conjecture 1.2 (Conjecture 29)**.** *For any group* G *the ideal of the group–based model is generated in degree at most* $|G|$ *for any tree* T*.*

Main Theorem 2. *For any group* G *there exists a constant* d *such that the projective scheme* $X_{\mathbb{P}}(T, G)$ *can be defined by an ideal generated in degree at most* d *for any tree* T*.*

The result presented above is quite similar to the main results of [DE], that generalize [DK09]. Let us state 3 major differences:

(1) The results in [DE] concern a wide class of *abelian equivariant* models, while ours deal only with *group–based* models;
(2) Our results are *scheme theoretic*, while the results in [DE] are *set theoretic*;
(3) Our results are *constructive*, while methods of [DE] are *existential.*

In [DE] the authors use a beautiful technique that basis on noetherian arguments in rings, that are not finitely generated, but are equipped with an additional monoid action. For more on this technique see [HS09, HdC09, Dra10]. Our methods are typical for toric geometry. Indeed, we believe that the presented approach can be used for much broader class of problems in toric geometry to determine the degree of generation – see for example [Bru11]. In particular the methods should cover group–based models for small groups G. As our proves are constructive, for group–based models, one can explicitly write a polynomial algorithm for testing if a point belongs to $X(T, G)$. The existence of such an algorithm was proven in [DE]. Still, the constants in general would be to big for the algorithm to be of practical value. However, for the 3-Kimura model, our results are sharp. The elements of the ideal defining $X(T, G)$ are called phylogenetic invariants. They are sought by people dealing with phylogenetics.

Corollary 1.3. *For any tree* T *any point* P *belongs to* $X(T, \mathbb{Z}_2 \times \mathbb{Z}_2)$ *if and only if all phylogenetic invariants of degree at most four vanish at* P*.*

The methods, how to inductively find phylogenetic invariants of a fixed degree were presented in [DBM, Section 3.2]. We believe that Corollary 1.3 may be used in applied mathematics.

In Section 2 we present a toric description of varieties $X(T, G)$. The general method that we use can be applied to other problems in toric geometry. It is described in Section 4. In Section 5 we present a basic combinatorial lemma. It is the main tool to prove Main Theorem 2, what we do in Section 6. The reader may notice the resemblance between *contractions* of tensors and methods of this section. The technical proof of Main Theorem 1 can be found in Section 7. In the last Section 8 we present open problems in the language of algebraic geometry.

Acknowledgements

The article was started at Universite Grenoble I and finished during a visit at Max Planck Institut in Bonn. I am grateful for their hospitality, especially to Laurent Manivel. I would like to thank Jarosław Wiśniewski for introducing me to the subject.

2. Constructions and notation

Let T be a rooted tree and G a finite abelian group. We assume that all edges of T are directed away from the root, however we could consider any orientation.

Definition 2.1 (Edges E, Vertices V, Leaves L, Nodes N)**.** *We define the sets E and V to be respectively the set of edges and vertieces of the tree T. Vertices of degree one are called leaves. The set of leaves is denoted by L. By abuse of notation, edges adjacent to leaves will also be called leaves. We define the set of nodes $N := V \setminus L$.*

There are two canonical groups one can consider.

Definition 2.2 (Edge labellings G_E, Vertex labellings G_V, Nodes labellings G_N)**.** *The group G_E of edge labellings consists of all functions $f : E \to G$. We define $(f_1f_2)(e) = f_1(e)f_2(e)$. The group G_E is isomorphic to $G^{|E|}$.*

The group G_V of vertex labellings consists of all functions $g : V \to G$. We define $(g_1g_2)(v) = g_1(v)g_2(v)$. The group G_V is isomorphic to $G^{|V|}$.

The group G_N of vertex labellings consists of all functions $g : N \to G$. We define $(g_1g_2)(v) = g_1(v)g_2(v)$. The group G_N is isomorphic to $G^{|N|}$.

As the tree T is directed there is a canonical morphism $s : G_E \to G_V$.

Definition 2.3 (The summing morphism s)**.** *We define $s : G_E \to G_V$ by $s(f)(v) = \sum_{e=(x,v)} f(e) - \sum_{e=(v,x)} f(e)$, where the first sum is taken over all edges incoming to v and the second over outgoing edges.*

In phylogenetics it is natural to distinguish the set of leaves and nodes. We define the natural projection π_N.

Definition 2.4 (Projection π_N)**.** *We define the projection $\pi_N : G_V \to G_N$ by restricting the domain of a function from V to N.*

The most important combinatorial objects that we consider are the following.

Definition 2.5 (Group-based flows)**.** *We define the group $\mathcal{G}$ to be the kernel of the composition $\pi_n \circ s : G_E \to G_N$. Hence the elements of $\mathcal{G}$ are such edge labellings that "the signed sum around each node is trivial". This is a well-known condition for a flow, but instead of labelling edges with numbers, we label them with group elements. We call elements of $\mathcal{G}$ group-based flows. We have got an exact sequence:*

$$1 \to \mathcal{G} \to G_E \to G_N \to 1.$$

As an abstract group we have an isomorphism $\mathcal{G} \cong G^{|L|-1}$.

Notice that in the definition above the role of sinks and sources is played by leaves. This construction can easily be generalized to other subsets of vertices.

Example 2.6. Let us consider the group $G = \mathbb{Z}_3$ and the following tree:



Here e_2, e_3, e_4 and e_5 are leaves. An example of a flow is an association $e_1 \to 2$, $e_2 \to 1$, $e_3 \to 1$, $e_4 \to 2$, $e_5 \to 2$.

Remark 2.7. Eventually we will see, Definition 2.12 and 2.15, that group-based flows correspond to vertices of the polytope associated to the toric variety. Such objects were first introduced only for the group $\mathbb{Z}_2$ and trivalent trees in [BW07, Definition 3.1] and were called networks. They were originally defined as pairwise disjoint paths, beginning and ending at leaves. Definition 2.5 is a direct generalization. One can associate to edges from paths an element $1 \in \mathbb{Z}_2$ and to all other edges $0 \in \mathbb{Z}_2$ obtaining a flow from a network.



It is worth noting, that as each element of the group in this case has order two, the definition of the flow is independent from the orientation of the tree.

It is well-known that, if the graph is a tree, it is enough to know the values at sources and sinks to reconstruct the flow. Of course, one cannot take arbitrary values for sources and sinks. For any flow, the sum over sources and sinks has to be equal to zero.

Definition 2.8 (Sockets $\mathcal{S}$)**.** *We define the group of sockets $\mathcal{S}$ whose elements are such functions $f : L \to G$ that $\sum_{l \in L} f(l) = e_0$, where e_0 is the neutral element of the group.*

Notice that by restricting a flow to leaves one obtains a socket. In fact one can easily prove the following.

Proposition 2.9. *The groups $\mathcal{G}$ and $\mathcal{S}$ are naturally isomorphic.* □

Remark 2.10. As it was for flows also sockets were first defined in [BW07, Definition 3.1] for trivalent trees and the group $\mathbb{Z}_2$. They were defined as even subsets of leaves. We see that the condition that the subsets are even corresponds to the summation condition in Definition 2.8.

We come to the definition of the polytope P that represents the variety $X_{\mathbb{P}}(T, G)$. This will be a lattice polytope in the lattice M_E.

Definition 2.11 (Lattice M_E, basis $b_{(e,g)}$)**.** *We define M_E as the lattice with basis elements indexed by all pairs $(e, g) \in E \times G$. The basis element indexed by (e, g) is denoted by $b_{(e,g)} \in M_E$.*

Definition 2.12 (Polytope P)**.** *To a flow $f \in \mathcal{G}$ one can naturally associate an element $P_f := \sum_{e \in E} b_{(e,f(e))} \in M_E$. This is an element whose coordinates are equal to either 1 or 0.*

We define P to be the convex hull of all the points P_f over all group-based flows f.

The following proposition follows from the fact that P is a subpolytope of the unite cube in M_E.

Proposition 2.13. *All integral points of P are vertices and are of the form P_f for certain flow $f \in \mathcal{G}$.* □

Definition 2.14 (Semigroup $S(P)$, Semigroup algebra $\mathbb{C}[P]$)**.** *We define the monoid $S(P)$ as the submonoid of M_E spanned by vertices of P. We define $\mathbb{C}[P]$ as the semigroup algebra associated to the monoid $S(P)$. Precisely $\mathbb{C}[P]$ is a complex vector space with basis elements identified with elements of $S(P)$. The multiplicative structure on $\mathbb{C}[P]$ is induced from the additive structure on $S(P)$.*

Definition 2.15 (Variety $X(T, G)$)**.** *Notice that $\mathbb{C}[P]$ is a graded algebra, taking vertices of P to be of degree 1. We define the projective variety $X_{\mathbb{P}}(T, G)$ as $\operatorname{Proj}\mathbb{C}[P]$. We also define $X(T, G) = \operatorname{Spec}\mathbb{C}[P]$.*

It is not obvious that the variety $X(T, G)$ agrees with the one defined in the Introduction. Indeed, it is a nontrivial theorem – [SS05, Mic11].

Remark 2.16. The variety $X_{\mathbb{P}}(T, G)$ is *not* the toric variety associated to the polytope P as defined for example in [Ful93, Section 1.5]. The variety $X_{\mathbb{P}}(T, G)$ does not have to be normal – for example when $G = \mathbb{Z}_6$ [DBM, Computation 4.3]. In fact, the toric variety associated to the polytope P is the normalization of $X_{\mathbb{P}}(T, G)$. We will be mostly interested in properties of $X(T, G)$. As the construction of $X(T, G)$ is motivated by applications, there are not many results on the normalization, that we find an interesting object for further studies.

Consider the vector space V with basis elements indexed by vertices of P. There is a natural embedding of $X_{\mathbb{P}}(T, G)$ in the projective space $\mathbb{P}(V)$. The choice of the basis induces a toric structure on $\mathbb{P}(V)$. The projective space $\mathbb{P}(V)$ contains a dense algebraic torus $\mathbb{T}$. The intersection of $\mathbb{T}$ with $X_{\mathbb{P}}(T, G)$ is also an algebraic torus that acts, with a dense orbit, on $X_{\mathbb{P}}(T, G)$. We are interested in the ideal $I(T, G)$ that defines $X_{\mathbb{P}}(T, G)$ in $\mathbb{P}(V)$.

The following crucial fact is well known in toric geometry.

Proposition 2.17 ([Stu96] Lemma 4.1)**.** *Let A be any finite set in a lattice M. Define the corresponding ideal I_A as in* [Stu96, Chapter 4]*. The toric ideal I_A is spanned, as a vector space, by the set of binomials*

$$\prod x_i^{a_i} - \prod x_i^{b_i},$$

where $\sum a_i A_i = \sum b_i B_i$ encodes an integral relation between points $A_i, B_i \in A$. □

We apply this proposition to $A = P$. We see that the description of the ideal $I(T, G)$ is *reduced to the study of integral relations between the vertices of the polytope P.*

2.1. **Group based polytopes.**

So far we have seen that the group $\mathcal{G}$ and polytope P are strongly related. We believe that the object consisting of such a pair deserves a separate, general definition.

Definition 2.18 (Group–based polytope)**.** *Let Q be a lattice polytope in the lattice M. Suppose that a group G acts on the integral points of Q. We say that the pair (Q, G) is a group based polytope if the G action preserves the linear relations. That is for any relation $\sum q_i = \sum q_j$ and any $g \in G$ we have $\sum g(q_i) = \sum g(q_j)$, where $q_i, q_j \in Q$ are lattice points.*

As before, the polytope Q generates a monoid $S(Q)$ which defines the semigroup algebra $\mathbb{C}[Q]$ -cf. Definition 2.14. The toric variety $\operatorname{Spec}\mathbb{C}[Q]$ has a natural embedding in the affine space V, with basis elements corresponding to lattice points of Q. As G acts on Q we see that V is a representation of G. The assertion in Definition 2.18 that the G action preserves the linear relation is equivalent to the fact that the action restricts to the variety $\operatorname{Spec}\mathbb{C}[Q]$.

It is straightforward to see that $(P, \mathcal{G})$ is a group based polytope. Moreover as the action of $\mathcal{G}$ on vertices of P is transitive and free we obtain the following proposition.

Proposition 2.19. *The natural ambient space V of the variety $\operatorname{Spec}\mathbb{C}[P]$ is the regular representation of the group $\mathcal{G}$. The induced action on $\mathbb{P}(V)$ restricts to the variety $X_{\mathbb{P}}(T, G)$.* □

Remark 2.20. In fact, canonically one should define flows as labellings of edges by *characters* of G, that is elements of G^*. As the groups G and G^* are isomorphic, in all constructions we get isomorphic objects. We decided not to introduce G^* to avoid unnecessary confusion.

3. General methods in phylogenetics

Although the construction from previous section takes several steps, still all the data that was used was the tree T and the group G. Thus one would expect to find certain relations between invariants of the ideal $I(G, T)$, the group G and the tree T.

Remark 3.1. An example of such a relation, in a slightly different setting, was presented in [BBKM10]. The genus of the graph plus one was proved to be the upper bound on the degree of generation of an algebra associated to any graph and the group $\mathbb{Z}_2$.

Definition 3.2 (Claw tree $K_{1,n}$)**.** *The claw tree $K_{1,n}$ is the tree with one node and n leaves.*

We introduce certain invariants that are of great interest [SS05, Chapter 5].

Definition 3.3 ($\phi(G, T), \phi(G), \phi(G, n)$)**.** *Let $\phi(G, T)$ be the (minimal) degree in which $I(G, T)$ is generated. Let $\phi(G, n) := \phi(G, K_{1,n})$. Let $\phi(G) = \sup \phi(G, T)$, where the supremum is taken over all trees.*

Let us restate the conjecture of Sturmfels and Sullivant.

Conjecture 3.4 ([SS05], Conjecture 29)**.** *We have $\phi(G) \leq |G|$.*

It was separately stated in the most interesting case for $G = \mathbb{Z}_2 \times \mathbb{Z}_2$, [SS05, Conjecture 30]. Note that, a priori, we do not know if $\phi(G)$ is finite, and indeed it is an open problem for groups different from $\mathbb{Z}_2$.

Let us present the combinatorial way of representing group-based flows and relations between vertices of P. Using the isomorphism $G_E \cong G^{|E|}$, each element of G_E can be represented as a sequence of group elements of length $|E|$. We represent group-based flows, as such sequences, presented as column vectors. A multiset of group-based flows is represented by a matrix. Each column corresponds to a flow and the rows are indexed by edges of the tree T. Recall that flows correspond to vertices of P. Consider any combination $\sum a_i v_i$ where $v_i \in P$ are vertices of P and $a_i \in \mathbb{N}$. Such a combination can be encoded as a matrix, where each flow corresponding to v_i appears a_i times. Of course such matrices should be considered up to permutation of columns.

Continuing this, we may encode a relation $\sum a_i v_i = \sum a_j v_j$ as a pair of matrices A_1, A_2 of the same size such that:

(1) rows are indexed by edges,
(2) each column represents a network,
(3) for each edge e the e-th row in A_1 is equal to the e-th row of A_2 up to permutation.

Example 3.5. Consider the following tree and numbering of edges:

(3.1)



Take $G = \mathbb{Z}_2$. An example of a nontrivial relation between flows is as follows:

$$\begin{bmatrix} 1 & 0 \\ 1 & 0 \\ 0 & 0 \\ 0 & 1 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 1 & 0 \\ 0 & 0 \\ 1 & 0 \\ 1 & 0 \end{bmatrix}.$$

One of the main methods in algebraic phylogenetics is reduction to simpler trees. Each tree T, different from the claw tree can be subdivieded into two simplier trees T_1 and T_2, by dividing an inner edge.



The operation can be reversed. If we have got two trees with distinguished leaves we can join them, obtaining one tree.

Often one can transfer certain results from the trees T_1 and T_2 to the tree T. This is the case for the degree of generation. We encourage the reader to proof the following result using the language of flows.

Proposition 3.6. *Suppose that we have got two trees T_1 and T_2 with distinguished leaves and that the tree T is obtained by joining them. Then* $\phi(G, T) \leq \max(\phi(G, T_1), \phi(G, T_2))$.□

In fact one can derive the generators of the ideal $I(G, T)$ from the ideal $I(G, T_1)$ and $I(G, T_2)$, [SS05, Theorem 26], [Sul07, Corollary 2.11]. Thus if we want to bound $I(G, T)$ for any tree T it is enough to consider claw trees.

Remark 3.7. The "reduction to claw trees" has been obtained in a much more general setting then presented in this paper [AR08, DK09]. Still the problem of describing the ideal for claw trees is extremely hard [DK09, p. 17], apart from the simple case $G = \mathbb{Z}_2$ [CP07].

In algebraic geometry varieties can be compared on different levels. The most common distinction is to compare them as *schemes* or as *sets*. Consider two ideals I_1, I_2 in a ring of polynomials $\mathbb{C}[x_1, \ldots, x_n]$. It is well-known that they define the same affine scheme if and only if they are equal. By Hilbert's Nullstellensatz we know that they define the same set if and only if their radicals are equal. We call the first equality *ideal theoretic* and the second one *set theoretic*. If the ideals are homogenous they define not only affine schemes, but also projective schemes. Let $m = (x_1, \dots, x_n)$ be the so called irrelevant ideal. The homogenous ideals I_1 and I_2 define the same projective scheme if and only if their saturation with respect to the maximal ideal are equal, that is $I_1 : m^\infty = I_2 : m^\infty$. Recall that $I : J^\infty = \{f : fJ^N \subset I \text{ for some } N \text{ large enough}\}$ is called saturation of I with respect to J. We call equality of ideals after saturation *scheme theoretic*. Of course equality of projective schemes implies equality of projective sets. So if both I_1 and I_2 are simultaneously either contained or not in m then *scheme theoretic* equality implies *set theoretic*.

Example 3.8. Consider $\mathbb{C}[x, y]$ and three ideals $I_1 = (x, y)$, $I_2 = (x^2, xy, y^2)$, $I_3 = (1)$. All of them define the same (empty) projective scheme. Their saturation is I_3. The first two define a nonempty affine set, supported at 0, with different scheme structure.

4. Idea

Definition 4.1 (Subideal I_d)**.** *For any graded ideal I we define the subideal generated by all elements of degree at most d by I_d.*

Let us start from general remarks on toric geometry. Consider a lattice polytope Q contained in a hyperplane H not passing through 0. This allows us to put a grading to the associated monoid generated by Q and hence on the associated toric ideal I_Q. Suppose that we want to prove that I_Q and I_{Qd} for some $d \in \mathbb{N}$ define the same projective scheme. Let X be the projective variety defined by Q. The integral points of Q correspond to coordinates of the ambient projective space of X. Proving that the saturation of I_{Qd} equals I_Q is equivalent to proving that I_{Qd} and I_Q are equal in each localization with respect to any coordinate, represented by a lattice point $R \in Q$. Thus we have to prove that any generator of I_Q multiplied by a sufficiently high power of the variable corresponding to R belongs to I_{Qd}.

Let us translate this condition to combinatorial language. The generators of I_Q correspond to relations between points of Q. Let us fix a relation $\sum A_i = \sum B_j$, where $A_i, B_j \in Q$. Multiplying the corresponding element of the ideal by the variable corresponding to R is equivalent to adding R to both sides of the relation. Thus we have to prove that the binomial corresponding to the relation $\sum A_i + mR = \sum B_j + mR$ is generated by binomials from I_Q of degree at most d for m sufficiently large.

A binomial corresponding to a relation $\sum R_i = \sum S_i$ between points of a polytope is generated in a degree d if and only if one can transform $\sum R_i$ to $\sum S_i$ using a sequence of simple steps. In each single transformation one can replace points $R_1, \dots, R_k$ for $k \le d$ by $R'_1, \dots, R'_k$ if they satisfy the relation $\sum_{i=1}^k R_i = \sum_{i=1}^k R'_i$. In such a case we say that the relation is generated in degree d.

The proof scheme is very simple:

(1) Using degree d relations reduce A_i, B_i to some simple, special points of Q contained in a subset $L \subset Q$. (*)
(2) Show that any relation between the points of L is generated in degree d.

In general any of this two points can be very difficult.

Remark 4.2. It is well known that the projective toric variety defined by a polytope Q is covered by affine subsets given by localizations by variables corresponding to vertices.

Thus one can be tempted to prove that $I_Q = I_{Qd}$ only in the localizations by variables corresponding to vertices. Note however that in general, we do not know if the scheme defined by I_{Qd} is also covered by open sets given by localizations by variables corresponding to vertices. Indeed, I_{Qd} and I_Q may be different on the set-theoretical level. For example if $\operatorname{Proj} I_{Qd}$ contains a point that is zero on the coordinates corresponding to vertices and nonzero on some other coordinates, then such a point will not belong to any open set corresponding to a localization with respect to vertices. However, if $\operatorname{Rad} I_{Qd} = I_Q$, then of course it is enough to consider localizations with respect to vertices. See also [Bru11].

As our polytopes P have only vertices, the problem described in Remark 4.2 does not concern us. We have the following equivalences for a toric ideal I_Q given by a polytope Q.

- All relations between vertices of Q are generated in degree $d \Leftrightarrow$ the ideal I is generated in degree d.
- For any point $R \in Q$ and any relation there is an integer m such that after adding mR to both sides of the relation, it is generated in degree $d \Leftrightarrow$ the projective scheme defined by I_Q can be defined by an ideal generated in degree d.

Lemma 4.3. *Suppose that P is a group–based polytope with G acting transitively on its integral points. The projective scheme $\operatorname{Proj}\mathbb{C}[P]$ can be represented by an ideal generated in degree at most d if and only if* there exists *a point $R \in P$ such that for any relation $\sum A_i = \sum B_i$ between points of P for m sufficiently large $\sum A_i + mR = \sum B_i + mR$ is generated in degree at most d.*

Proof. As G acts transitively the assumption that there exists a point R with the given property is equivalent to the fact that the property holds for all integral points of P. □

5. Combinatorial lemma

Definition 5.1 ([n])**.** *For any $n \in \mathbb{N}$ let $[n]$ be the set of natural numbers from zero to n.*

Definition 5.2 (Coloring)**.** *A coloring of length n is a function $f : [n] \to [i]$. The number i is called the number of colors. The support of the coloring f is defined as $\{k \in [n] : f(k) \neq 0\}$.*

Definition 5.3 (Transformation)**.** *Consider two colorings $f_1, f_2 : [n] \to [i]$. Suppose that there exist two numbers $0 \leq k_1, k_2 \leq n$ such that k_j is* not *in the support of f_j. Moreover suppose $f_1(k_2) = f_2(k_1)$. Define $f'_j(x) = f_j(x)$ for $x \neq k_1, k_2$. Moreover $f'_1(k_i) := f_2(k_i)$ and $f'_2(k_i) := f_1(k_i)$. We call f'_1, f'_2 a* transformation *of f_1 and f_2.*

Transformation corresponds to exchanging the fixed color in two colorings with a "blank" color. A set of colorings can be transformed into another by choosing two colorings and transforming them. We generate an equivalence relation on colorings by transformations. Abusing the notation the relation is also called transformation.

Lemma 5.4. *Let us fix two natural numbers i (number of colors) and s (bound on the support). Fix $\epsilon > 0$. There exists $N \in \mathbb{N}$, such that for all $n \geq N$ any collection of colorings $f_1, \ldots, f_m : [n] \to [i]$ with support of cardinality at most s can be transformed into a collection $f'_1, \ldots, f'_m$ such that there exist $(1-\epsilon)[\frac{n}{2}]$ even numbers $x < n$, such that for any f'_i at least one of the numbers $x, x+1$ is not in its support.*

Note that the statement is nontrivial, as the number m of colorings is not bounded.

Proof. The proof is inductive on s. For $s = 1$ we can take all even numbers smaller then n, thus the statement is trivial.

Assume that the statement is true for s. We will prove it for $s+1$, proceeding inductively on g. Fix ϵ.

Suppose $g = 1$. By induction hypothesis consider N' for s, $g = 1$ and $\epsilon' = \epsilon/2$. Consider $N = k(N' + 1)$ for large k. We call a coloring *bad* if its support is entirely contained in some interval $< m(N' + 1), (m + 1)(N' + 1))$. If we have two bad colorings with supports in different intervals we can transform them to two colorings that are not bad. Thus we can assume that all bad colorings are contained in one interval. By the induction hypothesis, we can transform the colorings, so that in each other interval, we can choose $(1 - \epsilon/2)[\frac{N'}{2}]$ even numbers. We see that for k large enough the theorem holds.

The case for greater g is only slightly more complicated. By induction, we choose N' that holds for $\epsilon/(g + 2)$, both in case when g *or* s is smaller. Consider $N = k(N' + 1)$. We call a coloring *bad* if both:

- its support is contained in some interval $< m(N' + 1), (m + 1)(N' + 1))$,
- all g colors appear in its image.

As before we can assume that all bad colorings are contained in one interval, say the first one. For any other interval, all colorings that are not bad can be divided into (not necessarily disjoint) subsets:

- g subsets depending on the color that is missing in the image,
- one subset containing colorings with supports not contained in the given interval.

For each coloring we choose one subset in which it is contained and fix this choice. By induction, for each interval apart from the first one, for each subset of colorings, there are at most $(\epsilon/(g + 2))[\frac{N}{2}]$ even numbers that do not satisfy the theorem. Hence, the theorem holds for k large enough. □

6. Bounding the degree of phylogenetic invariants

Definition 6.1 (Support of a flow)**.** *Let n be any flow. The set of edges to which n associates a nonneutral element is called the* support *of n.*

Lemma 6.2. *Consider a claw tree $K_{1,n}$ and a finite abelian group G. Let $\mathfrak{nt}$ be the neutral element in the group of flows. For any relation $\sum A_i = \sum B_i$ of integral points of the associated polytope P there exists m such that the relation $m\mathfrak{nt} + \sum A_i = m\mathfrak{nt} + \sum B_i$ can be transformed, using only quadrics, to a relation among flows with support at most $|G| - 1$.*

Proof. Consider any flow A. If its support is of cardinality greater that $|G| - 1$ we can find a strict subset S of edges such that $\sum_{e\in S} A(e)$ is the neutral element of G, where the addition is taken in G. Thus $\mathfrak{nt} + A$ equals the sum of two flows with strictly smaller support. The lemma follows easily. □

Lemma 6.3. *For any finite, abelian group G there exists N, such that if the ideal $I(K_{1,N}, G)$ is generated in degree d, then the scheme $X(K_{1,m}, G)$ can be defined by an ideal generated in degree d for any m.*

Proof. Let us choose N from Lemma 5.4 for $i = s = |G| - 1$ and $\epsilon < 1/2$. First let us prove the following claim:

Any relation between flows of length m with support of cardinality at most $|G| - 1$ is generated in degree d.

Let us prove it inductively on m, starting from $m = N$, where the assumption holds. Note that each flow can be considered as a coloring; we number group elements from 0 to $|G|-1$, assigning zero to the neutral element. As $\epsilon < 1/2$ we can find an even number x, such that x or $x+1$ is not in the support for any flow appearing in the relation. Let us replace each flow f appearing in the relation by:

$$f'(j) = \begin{cases} f(j) & j < x \\ f(j+1) & j > x \end{cases} \qquad f'(x) = \begin{cases} f(x) \text{ if } f(x) \neq e \\ f(x+1) \text{ if } f(x+1) \neq e \\ e \text{ if } f(x) = f(x+1) = e. \end{cases}$$

We obtain a relation between flows f' of length $m-1$. Thus, by induction, it is generated in degree d. Note that each generation step can be lifted to the relation between f, what proves the claim.

By multiplying any relation by the flow $\mathfrak{nt}$ in a sufficiently high power one can reduce any relation to relation between flows with support of cardinality at most $|G|-1$ using quadrics, Lemma 6.2. As flows act transitively on P, by Lemma 4.3 it is enough to consider relations multiplied by a high power of $\mathfrak{nt}$. Thus the lemma follows. □

Remark 6.4. *The constant N can be explicitly computed. Moreover due to standard theorems, a toric variety in bounded dimension and bounded exponents of defining monomials has got an explicitly bounded degree of generation* - [Stu96, Theorem 4.7, Corollary 4.15]. *Thus Main Theorem 2 follows.*

7. PROOF OF THE MAIN THEOREM 1

The whole Section is devoted to the proof of Main Theorem 1.

Theorem 7.1. *For any tree T the ideal $I(T, \mathbb{Z}_2 \times \mathbb{Z}_2)$ and the subideal generated in degree at most four define the same projective scheme.*

Following the arguments of [SS05, Chapter 5] one immediately reduces to the case when $T = K_{1,l}$ for certain $l \in \mathbb{N}$. Let us restate the general definitions already introduced, in case of the group $G = \mathbb{Z}_2 \times \mathbb{Z}_2$ (the 3-Kimura model) on a claw tree $K_{1,l}$.

Definition 7.2 (Flow, ideals I and I_4, polytope P)**.** *A flow is an association of elements of the group $\mathbb{Z}_2 \times \mathbb{Z}_2$ to edges of $K_{1,l}$, such that the sum of all elements is the natural element. Let I be the ideal of the variety $X(K_{1,l}, \mathbb{Z}_2 \times \mathbb{Z}_2)$ and let I_4 be the subideal generated in degree 4. Let P be the corresponding polytope.*

We will identify a flow with an l-tuple of group elements summing up to zero. The sum of such l-tuples will be a coordinatewise sum, where each entry is treated as an element of the free abelian group generated by elements of $\mathbb{Z}_2 \times \mathbb{Z}_2$. Each flow represents a vertex of a polytope P. The addition described above is the addition in the lattice generated by vertices of the polytope.

Example 7.3. For $l = 4$ we can add:

$$(0, (1,0) + (1,1), 2(0,1), -3(0,0)) + ((0,1), (1,0) - (1,1), (1,0), (0,0))$$
$$= ((0,1), 2(1,0), 2(0,1) + (1,0), -2(0,0)).$$

The trivial flow is $((0,0), (0,0), (0,0), (0,0))$.

Definition 7.4 (Pair, triple)**.** *We say that a flow is a* pair *if and only if the cardinality of the support is equal to two. We say that a flow is a* triple *if and only if the cardinality of the support is equal to three.*

By Lemmas 4.3 and 6.2 we have to generate relations only between flows that are pairs and triples. This completes the first step of the method (*) presented in Section 4. The set L consists of pairs and triples. Note that this part of the proof can be adjusted to other groups G, for L consisting of flows with support of cardinality at most $|G|-1$.

Let us fix any relation $\sum n_i = \sum n'_i$, where n_i and n'_i are either pairs or triples. Our aim is to transform $\sum n_i$ to $\sum n'_i$ in a series of steps, each time replacing at most four n_i by flows with the same sum[2]. We assume that among n_i there are more or the same number of triples as among n'_i. We first try to reduce the relation, so that consequently:

1. Among n_i there are as few triples as possible,
2. Among n'_i there are as few triples as possible,
3. The degree of the relation is as small as possible.

More precisely let t and t' be the number of triples among respectively n_i and n'_i. Let d be the degree of the relation. Our proof will be inductive on (t, t', d) with lexicographic order.

To prove Theorem 7.1 we consider separately three cases depending on the number of triples among n_i. The cases are:

a) there are no triples,
b) there is exactly one triple,
c) there are at least two triples.

We say that a family of flows agrees on an index j of an edge if they all associate the same element to j and j belongs to their *support*. We will denote by g_1, g_2 and g_3 the three nontrivial elements of $\mathbb{Z}_2 \times \mathbb{Z}_2$. A triple that associates g_1 to index a, g_2 to index b and g_3 to index c is denoted by (a, b, c). A pair that associates an element g_i to indices d and e will be denoted by $(d, e)_{g_i}$. We say that g_i is contained in a flow if there exists an index j, such that the flow associates g_i to j. We believe that the following proofs are impossible to follow without a piece of paper. We strongly encourage the reader to note what flows appear on both sides of the relation at each step of the proof.

7.1. **The case with no triples.** First note that there are no triples among n'_i. Without loss of generality we may assume that n_1 is a pair equal to $(a, b)_{g_1}$. Hence there exists n'_i, say n'_1, that is $(b, c)_{g_1}$ for some index c. If $c = a$ we can reduce this pair, hence we assume $c \neq a$. There exists a flow, say n_2 that is $(c, d)_{g_1}$. If $d = b$ we can reduce this pair. We consider two other cases:

1) $d \neq a$. Then we use the degree two relation $(a, b)_{g_1} + (c, d)_{g_1} = (a, d)_{g_1} + (b, c)_{g_1}$ and we can reduce $(b, c)_{g_1}$.

2) $d = a$. Then there is a flow, say n'_2 given by $(a, e)_{g_1}$. If $e = b$ or $e = c$ we can reduce this pair. In the other cases we use the relation $(a, e)_{g_1} + (b, c)_{g_1} = (a, b)_{g_1} + (e, c)_{g_1}$ and we reduce $(a, b)_{g_1}$.

Notice that in this very easy case we have only used degree two relations.

7.2. **The case with one triple.** Let n_1 be the only triple among n_i.

Lemma 7.5. *There is exactly one triple among n'_i.*

---

[2] We are also allowed to add the flow $\mathfrak{nt}$ to both sides.

Proof. Due to the assumptions we know that there is at most one triple among n_i'. We exclude the case when there are no triples by comparing the parity of the number of times the element g_1 appears on both sides of the relation. ☐

Due to the previous lemma we may assume that n_1' is the only triple among n_i'. Let $n_1 = (1, 2, 3)$.

7.2.1. *Case: the triples agree on at least two elements in their support.* Consider the case where triples agree on two elements, say 1 and 2, that are in their support. Suppose that $n_1 = (1, 2, 3)$ and $n_1' = (1, 2, c)$. Of course if $c = 3$ we can make a reduction. In other case we must have a pair $(c, d)_{g_3}$ among n_i. If $d \neq 3$ then we use the relation $(c, d)_{g_3} + (1, 2, 3) = (1, 2, c) + (3, d)_{g_3}$ and reduce the triples. Hence $d = 3$. Analogously there must be a pair $(3, c)_{g_3}$ among n_i', hence we can reduce this pair.

7.2.2. *Case: the triples agree on exactly one element in their support.* Consider the case when triples agree on at least one element, say 1, in their common support. By the previous case we may assume that they agree on exactly one element.

As before let $n_1 = (1, 2, 3)$ and $n_1' = (1, b, c)$. We consider three cases.

1) $b \neq 3$.

There must be a pair $(b, d)_{g_2}$ among n_i. If $d \neq 2$ then we can apply the relation $(b, d)_{g_2} + n_1 = (1, b, 3) + (d, 2)_{g_2}$. This reduces to the case 7.2.1. So we assume $d = 2$. There must be a pair $(2, e)_{g_2}$ among n_i'. Hence there must also be a pair $(e, f)_{g_2}$ among n_i. If $f \neq b$ we can use a relation $(e, f)_{g_2} + (2, b)_{g_2} = (e, 2)_{g_2} + (f, b)_{g_2}$ and reduce $(e, 2)_{g_2}$. For $f = b$ we must have a pair $(b, g)_{g_2}$ among n_i'. If $g = 2$ or $g = e$ then this pair can be reduced. In the other case we use the relation $(e, 2)_{g_2} + (b, g)_{g_2} = (e, g)_{g_2} + (b, 2)_{g_2}$ and reduce $(b, 2)_{g_2}$.

2) $c \neq 2$.

This case is analogous to 1).

3) $b = 3$ and $c = 2$.

Lemma 7.6. *If there is a pair $(p, q)_{g_2}$ among n_i, such that $p, q \neq 2$ then we may assume that it is equal to* $(1, 3)$.

Proof. Suppose that $p \neq 1, 3$ and $q \neq 2$. We apply a relation $(p, q)_{g_2} + n_1 = (1, p, 3) + (q, 2)_{g_2}$ and reduce to case 1) $b \neq 2$. ☐

Analogously if there is a pair $(p, q)_{g_2}$ among n_i', such that $p, q \neq 3$ then this pair equals $(1, 2)_{g_2}$.

Notice that there must be a pair $(3, d)_{g_2}$ among n_i and a pair $(2, e)_{g_2}$ among n_i'. From Lemma 7.6 d equals 2 or 1 and e equals 3 or 1. We will consider subcases.

3.1) Suppose that $d = 2$.

If $e = 3$ then we can make a reduction of pairs. If $e = 1$ we must have a pair $(1, f)_{g_2}$ among n_i. If $f = 2$ we make a reduction, hence we assume $f = 3$. This means that there must be a pair $(3, g)_{g_2}$ among n_i'. If $g = 2$ or $g = 1$ we can make a reduction. Otherwise we apply the relation $(1, 2)_{g_2} + (3, g)_{g_2} = (1, 3)_{g_2} + (2, g)_{g_2}$ and reduce the pair $(1, 3)_{g_2}$.

3.2) Suppose that $e = 3$.

This case is similar to 3.1).

3.3) Suppose that $d = 1$ and $e = 1$.

As this is the only case left we may repeat the same reasoning for g_3. In particular, we must have a pair $(1, 2)_{g_3}$ among n_i. We see that we can reduce the triples by applying the following relation:

$$(1,2,3)+(1,3)_{g_2}+(1,2)_{g_3}=(1,3,2)+(1,2)_{g_2}+(1,3)_{g_3}.$$

This is a degree three relation.

7.2.3. *Case: The triples do not agree on any element of the support.* We want to reduce to one of previous cases. Consider the following two cases.

1) *The triples n_1 and n'_1 have different supports.*

Once again let $(1,2,3)=n_1$ and let $(a,b,c)=n'_1$. We may assume that a is not in the support of n_1. We see that there must be a pair $(a,f)_{g_1}$ among n_i. If $f\neq 1$ we can use a relation $(a,f)_{g_1}+n_1=(a,2,3)+(f,1)_{g_1}$. This reduces to the case 7.2.2, hence we assume that $f=1$. There must be a pair $(g,1)_{g_1}$ among n'_i. If $g=a$ we can reduce this pair, so we assume $g\neq a$. Notice that there must be a pair $(g,h)_{g_1}$ among n_i. If $h\neq a$, then we can use relation $(1,a)_{g_1}+(g,h)_{g_1}=(g,1)_{g_1}+(h,a)_{g_1}$ and reduce the pair $(g,1)_{g_1}$. So we can assume $h=a$. Then there must be a pair $(a,i)_{g_1}$ among n'_i. If $i=1$ then we can reduce it. Otherwise we can use the relation $(g,1)_{g_1}+(a,i)_{g_1}=(g,a)_{g_1}+(1,i)_{g_1}$ and reduce the pair $(g,a)_{g_1}$.

2) *The set $\{1,2,3\}$ is the support of n'_1 and n_1.*

Remember that due to the assumption 7.2.3 the triples n_1 and n'_1 do not agree on any element from their support. Without loss of generality we may assume $n_1=(1,2,3)$ and $n'_1=(2,3,1)$. Hence there must be a pair $(2,a)_{g_1}$ among n_i and $(1,b)_{g_1}$ among n'_i. If $a=1$ and $b=2$ then both pairs are the same and can be reduced. As both cases are symmetric we can assume that $a\neq 1$.

If $a\neq 3$ we can use the relation $(2,a)_{g_1}+(1,2,3)=(a,2,3)+(2,1)_{g_1}$. This reduces to the case with different supports. We are left with the case $a=3$. There must be a pair $(3,z)_{g_1}$ among n'_i. If $z\neq 1$ we can use the relation $(3,z)_{g_1}+n'_1=(z,3,1)+(2,3)_{g_1}$. This would enable to reduce the $(2,3)_{g_1}$ pair and decrease the degree. So we can assume that $z=1$. So far we have shown that there must be pairs $(2,3)_{g_1}$ among n_i and $(3,1)_{g_1}$ among n'_i[3]. By the same reasoning for g_2 and g_3 we see that we can use the following relation:

$$(1,2,3)+(2,3)_{g_1}+(1,3)_{g_2}+(1,2)_{g_3}=(2,3,1)+(2,3)_{g_3}+(1,3)_{g_1}+(1,2)_{g_2}.$$

Notice that this is a degree four relation. It enables us to reduce triples.

7.3. **The case with at least two triples.** We suppose that there are at least two triples among n_i.

Lemma 7.7. *If there are two triples n_1, n_2 among n_i that do not agree on any element of their supports then we can make a reduction. Thus we can assume that any two triples among n_i agree on at least one index.*

Proof. The assumptions are equivalent to $n_1=(a,b,c)$, $n_2=(d,e,f)$ with $a\neq d$, $b\neq e$, $c\neq f$. We apply the relation $n_1+n_2+\mathfrak{nt}=(a,d)_{g_1}+(b,e)_{g_2}+(c,f)_{g_3}$ that reduces the number of triples. ☐

Lemma 7.8. *If there is no index on which all triples from n_i agree then we can make a reduction.*

---

[3] Notice that we have made a symmetry assumption $a\neq 1$. The symmetric assumption would be $b\neq 2$. However as the result we got was symmetric, also for $b\neq 2$ we prove the existence of the same pairs.

Proof. Suppose there is no index on which all n_i agree. We may consider only two cases due to Lemma 7.7.

1) Suppose that any two triples from n_i agree on at least two elements.

Consider any triple $n_1 = (1, 2, 3)$. Due to the fact that not all triples from n_i associate g_1 to 1 there is a triple $(a, 2, 3)$ with $a \neq 1$ among n_i. There also must be a triple that does not associate g_2 to 2. It cannot agree both with $(1, 2, 3)$ and $(a, 2, 3)$ on two indices.

2) There exist two triples that agree only on one index.

Let $n_1 = (1, 2, 3)$ and $n_2 = (1, b, c)$ with $b \neq 2$ and $c \neq 3$. Due to the case assumption there is a triple $n_3 = (d, e, f)$ with $d \neq 1$. Remember that any two triples have to agree on at least one element due to Lemma 7.7. Hence without loss of generality we can assume $e = b$ and $f = 3$. We can apply the relation:

$$n_1 + n_2 + n_3 + \mathfrak{nt} = (d, 1)_{g_1} + (2, b)_{g_2} + (3, c)_{g_3} + (1, b, 3),$$

that reduces the number of triples. □

Due to the previous lemma we may assume that there exists an index, say 1*, such that all triples among* n_i *associate to it the same nonneutral element, say* g_1*.*

Definition 7.9 (k)**.** *Let* k *be the number of indices on which all triples among* n_i *agree. We know that* $1 \leq k \leq 3$*.*

We proceed inductively on k, as for $k = 0$ we already know how to reduce the relation. Hence from now on decreasing k is also a reduction.

Lemma 7.10. *Suppose that all triples from* n_i *associate* g_j *to an index* l*. If there is a pair* $(x, y)_{g_j}$ *among* n_i *with* $l \neq x, y$ *then either* $\{l, x, y\}$ *is the support of all triples among* n_i *or we can make a reduction.*

Proof. To simplify the language assume $g_j = g_1$ and $l = 1$. Suppose that there is a triple $n_1 = (1, b, c)$ with the support different from $\{1, x, y\}$. We can assume $x \neq b, c$. We apply the relation $n_1 + (x, y)_{g_1} = (x, b, c) + (1, y)_{g_1}$ what reduces k. □

Lemma 7.11. *Suppose that all triples from* n_i *associate* g_j *to an index* l*. If all pairs* $(x, y)_{g_j}$ *among* n_i *have* l *in the support then we can reduce all such pairs.*

Proof. Let t be the number of triples among n_i. Let p be the number of g_j pairs among n_i. Let t'_1 and t'_2 be the number of triples in n'_i that respectively assign or do not assign g_j to l. Let p'_1 and p'_2 be the number of g_j pairs among n'_i that respectively have or do not have l in the support. We know that $t \geq t'_1 + t'_2$. Comparing the number of times g_j appears in n_i and n'_i we get:

$$t + 2p = t'_1 + t'_2 + 2(p'_1 + p'_2).$$

Comparing the number of times g_j appears on index l we get:

$$t + p = t'_1 + p'_1.$$

This forces $t'_2 = p'_2 = 0$, $t = t'_1$ and $p = p'_1$. Hence all g_j pairs and triples among n_i and n'_i must assign g_j to l. Hence the multisets of pairs must be the same for n_i and n'_i. □

Lemma 7.12. *If there are* g_1 *pairs among* n_i*, then we can make a reduction.*

Proof. We will prove that there are no pairs $(a,b)_{g_1}$ among n_i that do not have 1 in the support. Due to Lemma 7.11 this will finish the proof. Suppose that there is a pair $(a,b)_{g_1}$ among n_i with $a,b \neq 1$. Due to Lemma 7.10 all the triples among n_i must have the support $\{1,a,b\}$. So either $k=1$ or $k=3$. If $k=1$ we can apply the relation

$$(1,a,b)+(1,b,a)+(a,b)_{g_1}+\mathfrak{nt}=(1,a)_{g_1}+(1,b)_{g_1}+(a,b)_{g_2}+(a,b)_{g_3}.$$

This reduces the number of triples. Thus we can assume that all triples among n_i are equal to $(1,a,b)$.

Claim: *Consider any pair $(c,d)_{g_2}$ among n_i. We can assume that its support is contained in $\{1,a,b\}$.*

Proof of the Claim. Suppose this is not the case, that is $c \notin \{1,a,b\}$. Due to Lemma 7.10 we can assume $d=a$.

1) Suppose that there is a g_2 pair among n_i that does not contain a in the support.

It must be equal to $(1,b)_{g_2}$ due to Lemma 7.10. We can apply the relation $(1,b)_{g_2}+(a,c)_{g_2}=(c,1)_{g_2}+(a,b)_{g_2}$. Applying once again Lemma 7.10 to the pair $(c,1)_{g_2}$ we can make a reduction.

2) All g_2 pairs among n_i contain a in the support.

Due to Lemma 7.11 we can make a reduction. ☐

Thus the support of all g_2 pairs among n_i is contained in $\{1,a,b\}$. The same holds for g_1 and g_3 pairs. Thus all flows among n_i have support contained in $\{1,a,b\}$. Hence the same must hold for n_i'. So our relation is a relation only on three indices. It is well known that the ideal for a tree with three edges is generated in degree 4, so in particular the considered relation is generated in degree 4. ☐

Corollary 7.13. *If all triples among n_i associate g_j to an index l, then there are no g_j pairs among n_i. Consequently there are no g_j pairs among n_i' and all triples among n_i' associate g_j to l. Moreover the number of triples among n_i equals the number of triples among n_i'.* ☐

We reduced to the case were there are no g_1 pairs neither among n_i nor n_i'. Moreover, there is the same number of triples among n_i and n_i' and they all associate g_1 to 1.

Lemma 7.14. *If all the triples among n_i and n_i' have support in $\{1,2,3\}$ then we can make a reduction.*

Proof. In this case $k=1$ or $k=3$. If $k=1$ then among n_i there is a triple $(1,2,3)$ and $(1,3,2)$. One of this triples can be reduced. If $k=3$ there are no pairs. All triples among n_i and n_i' are equal, thus the relation is trivial. ☐

7.3.1. *Case:* $k=1$. We first consider the most difficult case $k=1$. As always let $n_1=(1,2,3)$ and $n_1'=(1,b,c)$. As the proof is quite complicated we decided to include the tree that describes most important cases. While reading the proof we encourage the reader to follow at which node we are. The proof is "depth-first, left-first".



We start with the left node in the second row – assume $b=2$.

We move to the most left node in the third row – suppose that there is no g_3 pair among n_i that has got c in the support and, symmetrically, there is no g_3 pair among n_i' that has got 3 in the support. There must be a triple $(1,e,c)$ among n_i. If $e\neq 3$ then we apply the relation $(1,2,3)+(1,e,c)=(1,2,c)+(1,e,3)$ and reduce the triple $(1,2,c)$. We have $e=3$. Analogously among n_i' there must be a triple $(1,c,3)$. Hence there must be either a pair $(c,f)_{g_2}$ or a triple $(1,c,g)$ among n_i.

We continue to the most left node in the fourth row – suppose that there is a pair $(c,f)_{g_2}$. If $f\neq 2$ we apply the relation $(1,2,3)+(c,f)_{g_2}=(1,c,3)+(f,2)_{g_2}$ and reduce the triple $(1,c,3)$. If $f=2$ we apply the relation $(1,3,c)+(c,2)_{g_2}=(1,2,c)+(3,c)_{g_2}$ and reduce the triple $(1,2,c)$.

Hence we can assume that there is a triple $(1,c,g)$ among n_i – second node in the fourth row. If $g\neq 2$ then we apply the relation $(1,c,g)+(1,2,3)=(1,2,g)+(1,c,3)$ and reduce the triple $(1,c,3)$. For $g=2$ we apply the relation $(1,2,3)+(1,3,c)+(1,c,2)=(1,2,c)+(1,3,2)+(1,c,3)$ and reduce the triple $(1,2,c)$.

We continue to the second node in the third row. We assume that there is a pair $(3,l)_{g_3}$ among n_i'. If $l\neq c$ we apply the relation $(1,2,c)+(3,l)_{g_3}=(1,2,3)+(c,l)_{g_3}$ and reduce the triple $(1,2,3)$. If there was a pair $(c,m)_{g_3}$ among n_i then analogously we would have $m=3$ and we would be able to reduce this pair. So there must be a triple $(1,n,c)$ among n_i. If $n\neq 3$ then we apply the relation $(1,2,3)+(1,n,c)=(1,n,3)+(1,2,c)$ and reduce the triple $(1,2,c)$. So we assume $n_2=(1,3,c)$. Hence there is either a pair $(3,o)_{g_2}$ or a triple $(1,3,p)$ among n_i'.

We move to the third node in the fourth row – suppose that there is a triple $(1,3,p)$ among n_i'. If $p\neq 2$ we apply the relation $(1,2,c)+(1,3,p)=(1,2,p)+(1,3,c)$ and we reduce $(1,3,c)$. So we have $p=2$. We apply the relation $(1,3,2)+(3,c)_{g_3}=(1,3,c)+(2,3)_{g_3}$ and reduce the triple $(1,3,c)$.

We pass to the fourth node in the fourth row – we assume that there is a pair $(3,o)_{g_2}$ and there is no triple $(1,3,p)$ among n_i'. If $o\neq 2$ then we apply the relation $(1,2,c)+(3,o)_{g_2}=(1,3,c)+(2,o)_{g_2}$ and reduce $(1,3,c)$. So there is a pair $(2,3)_{g_2}$ among n_i'. Suppose that this pair appears $r>0$ times among n_i'. Note that there are no pairs $(2,s)_{g_2}$ among n_i. Indeed suppose that there is such a pair. If $s\neq 3$ then we apply the relation $(1,3,c)+(2,s)_{g_2}=(1,2,c)+(3,s)_{g_2}$ and reduce the triple $(1,2,c)$. If $s=3$ we reduce the pair $(2,3)_{g_2}$. Hence there must be at least $r+1$ triples of the type $(1,2,t)$ among n_i. If there is a triple with $t\neq 3$ then we apply the relation $(1,3,c)+(1,2,t)=(1,3,t)+(1,2,c)$ and reduce the triple $(1,2,c)$. Hence we have got at least $r+1$ triples $(1,2,3)$ among n_i. Notice that there are no triples of the type $(1,y,3)$ among n_i'. Indeed, in such a case we could apply the relation $(1,y,3)+(2,3)_{g_2}=(1,2,3)+(y,3)_{g_2}$ and reduce $(1,2,3)$. Hence there must be at least $r+1$ pairs of the type $(3,u)_{g_3}$ among n_i'. If $u\neq c$ then we apply the relation $(1,2,c)+(3,u)_{g_3}=(1,2,3)+(c,u)_{g_3}$ and reduce the triple $(1,2,3)$. Hence we have at least $r+1$ pairs $(3,c)_{g_3}$ among n_i'. Note that there are no pairs of the type $(c,v)_{g_3}$ among n_i. Indeed if $v=3$ we could reduce this pair. If $v\neq 3$ then we apply the relation $(1,2,3)+(c,v)_{g_3}=(1,2,c)+(3,v)_{g_3}$ and reduce the triple $(1,2,c)$. Hence we must have at least $r+1$ triples of the type $(1,z,c)$ among n_i. If $z\neq 3$ then we apply the relation $(1,2,3)+(1,z,c)=(1,2,c)+(1,z,3)$ and reduce the triple $(1,2,c)$. So there are at least $r+1$ triples $(1,3,c)$ among n_i. Note that the elements g_2 on 3 cannot be reduced – among n_i' there are only r pairs containing them and no triples. The contradiction finishes this case.

Consider the third node in the third row – there is a pair $(c,w)_{g_3}$ among n_i. This is completely analogous to the second node in this row, already considered.

Also the second node in the second row – $c=3$ – is analogous to the first node in the second row.

We are left with the last, third node in the second column – any two triples n_i and n_j' agree on exactly one index, that is on 1. Due to Lemma 7.14 we can assume $b\neq 2$ and $b\neq 3$. Due to the case assumption there must be a pair $(b,d)_{g_2}$ among n_i. If $d\neq 2$ then we apply the relation $(1,2,3)+(b,d)_{g_2}=(1,b,3)+(d,2)_{g_2}$ and reduce to the case $b=2$[4]. Analogously we must have the same pair among n_i' and it can be reduced.

7.3.2. *Case: $k=2$ or $k=3$.* Suppose now that $k=2$. Let $n_1=(1,2,3)$ and $n_1'=(1,2,c)$. If we cannot reduce n_1' then there must be a pair $(c,d)_{g_3}$ among n_i and a pair $(3,e)_{g_3}$ among n_i'. If $d=3$ and $e=c$ we can reduce the pairs. Thus we can assume that $d\neq 3$. We apply the relation $(1,2,3)+(c,d)_{g_3}=(1,2,c)+(3,d)_{g_3}$ and reduce the triple $(1,2,c)$.

The last, easiest case is $k=3$. Then all triples are equal to $(1,2,3)$ and there are no pairs due to Corollary 7.13. Hence we can reduce the triples. This finishes the proof of the theorem.

8. Open problems

We have already recalled two conjectures of Sturmfels and Sullivant 1.1 and 1.2. One can state simpler conjectures, that we find important.

Conjecture 8.1. *For any group G and any tree T the ideal $I(X(G,T))$ and the subideal generated in degree $|G|$ define the same projective scheme.*

We believe that the methods of Section 7 could be adopted, especially for groups of small cardinality.

Consider a claw tree $K_{1,n}$. Subdivide the set of leaves into two separate subsets. By adding an inner edge that separates this set one obtains a tree T as shown below.



---

[4] Notice that we do not reduce to the case $k=2$ as if this was true we would have already been in the first node in the second column $b=2$.

One can easily see that for any[5] group G we have $X(K_{1,n}, G) \subset X(T, G)$. As long as the subsets of leaves are of cardinality at least 2 the procedure decreases the maximal degree and allows for an inductive generation of phylogenetic invariants.

Conjecture 8.2 ([DBM] Conjecture 3.6)**.** *The variety $X(K_{1,n}, G)$ is a scheme theoretic intersection of all $X(T, G)$, where T is a prolongation with maximal degree at most $n-1$.*

This conjecture would imply the following.

Conjecture 8.3. *The function $\phi(G, \cdot)$ is constant.*

In view of the previous conjecture and Conjecture 1.2 it is natural to bound $\phi(G, K_{1,3})$. It is an open problem if $\phi(G, K_{1,3}) \leq |G|$. On the other hand, by general [Stu96, Theorem 4.7] we have an exponential bound on $\phi(G, K_{1,3})$. The following question, as far as we know, is open and would be an interesting step towards conjectures of Sturmfels and Sullivant.

Conjecture 8.4. *There exists a polynomial P such that $\phi(G, K_{1,3}) \leq P(G)$.*

One obtains a much more difficult open problem by replacing 3 in the previous conjecture by any (fixed or even non fixed) number.

The following conjecture concerns normality of varieties $X(T, G)$.

Conjecture 8.5. *The variety $X(T, \mathbb{Z}_2 \times \mathbb{Z}_2)$ is projectively normal.*

The previous conjecture is known for trees of small maximal degree. Note that $X(K_{1,3}, \mathbb{Z}_6)$ is not normal [DBM, Computation 4.3]. However analogous questions for many other groups are open. For the group $G = \mathbb{Z}_2$ the reader can consult [DBM].

For group–based models the cardinality of the group G equals the dimension of the vector space V associated to any node. The dimension of the space associated to the node, for the general Markov model determines which secant of the Segre we consider. It is well–known (and easy to prove) that there are no equations of degree less or equal to k for the k-th secant.

Conjecture 8.6. *Is the k-th secant of the Segre variety defined by equations of degree $f(k)$ where f is a linear function?*

References

[AR08] Elizabeth S. Allman and John A. Rhodes, *Phylogenetic ideals and varieties for the general Markov model*, Advances in Applied Mathematics **40(2)** (2008), 127–148.

[BBKM10] Weronika Buczyńska, Jarosław Buczyński, Kaie Kubjas, and Mateusz Michalek, *Degrees of generators of phylogenetic semigroups on graphs*, arXiv:1004.1183v1 [math.AG] (2010).

[Bru11] Winfried Bruns, *The quest for counterexamples in toric geometry*, Preprint at arXiv:1110.1840 (2011).

[BW07] Weronika Buczyńska and Jarosław A. Wiśniewski, *On geometry of binary symmetric models of phylogenetic trees*, J. Eur. Math. Soc. **9(3)** (2007), 609–635.

[CP07] J. Chifman and S. Petrović, *Toric ideals of phylogenetic invariants for the general group-based model on claw trees $k_{1,n}$*, Proceedings of the 2nd international conference on Algebraic biology (2007), 307–321.

[DBM] Maria Donten-Bury and Mateusz Michałek, *Phylogenetic invariants for group-based models*, arXiv:1011.3236v1.

[DE] J. Draisma and R.H. Eggermont, *Finiteness results for abelian tree models*, arXiv:1207.1282v1.

---

[5] In fact for any model.

[DK09] Jan Draisma and Jochen Kuttler, *On the ideals of equivariant tree models*, Mathematische Annalen **344(3)** (2009), 619–644.

[DK11] ______, *Bounded-rank tensors are defined in bounded degree*, arXiv:1103.5336v2 (2011).

[Dra10] Jan Draisma, *Finiteness for the k-factor model and chirality varieties*, Adv. Math. **223** (2010), no. 1, 243–256. MR 2563217 (2011e:14091)

[ERSS04] N. Eriksson, K. Ranestad, B. Sturmfels, and S. Sullivant, *Phylogenetic algebraic geometry*, Projective Varieties with Unexpected Properties; Siena, Italy (2004), 237–256.

[ES93] Steven N. Evans and T. P. Speed, *Invariants of some probability models used in phylogenetic inference*, Ann. Statist. **21(1)** (1993), 355–377.

[Ful93] William Fulton, *Introduction to toric varieties*, Annals of Mathematics Studies, vol. 131, Princeton University Press, Princeton, NJ, 1993, The William H. Roever Lectures in Geometry.

[HdC09] Christopher J. Hillar and Abraham Martin del Campo, *Finiteness theorems and algorithms for permutation invariant chains of laurent lattice ideals*, arXiv:1110.0785 (2009).

[HP89] Michael D. Hendy and David Penny, *A framework for the quantitative study of evolutionary trees*, Systematic Zoology **38** (1989), 297–309.

[HS09] Christopher J. Hillar and Seth Sullivant, *Finite groebner bases in infinite dimensional polynomial rings and applications*, arXiv:0908.1777 (2009).

[JC69] T. H. Jukes and C. R. Cantor, *Evolution of protein molecules*, Mammalian protein metabolism (M. N. Munro, ed.), vol. III, Academic Press, N. Y., 1969, pp. 21–132.

[Kim80] M. Kimura, *A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences*, Journal of Molecular Evolution **16** (1980), 111–120.

[Kim81] ______, *Estimation of evolutionary distances between homologous nucleotide sequences*, Proceedings of the National Academy of Sciences **78** (1981), no. 1, 454–458.

[Man09] C. Manon, *The algebra of Conformal Blocks*, arXiv:0910.0577 (2009).

[Mic11] Mateusz Michałek, *Geometry of phylogenetic group-based models*, Journal of Algebra **339** (2011), 339–356.

[PS05] Lior Pachter and Bernd Sturmfels (eds.), *Algebraic statistics for computational biology*, Cambridge University Press, New York, 2005.

[Rai11] Claudiu Raicu, *Secant varieties of segre–veronese varieties*, arXiv:1011.5867v2 (2011).

[SS03] C. Semple and M. Steel, *Phylogenetics*, Oxford University Press, 2003.

[SS05] Bernd Sturmfels and Seth Sullivant, *Toric ideals of phylogenetic invariants*, J. Comput. Biology **12** (2005), 204–228.

[SSE93] L. A. Székely, M. A. Steel, and P. L. Erdös, *Fourier calculus on evolutionary trees*, Appl. Math. **14(2)** (1993), 200–210.

[Stu96] Bernd Sturmfels, *Gröbner bases and convex polytopes*, University Lecture Series, vol. 8, American Mathematical Society, 1996.

[Sul07] Seth Sullivant, *Toric fiber products*, Journal of Algebra **316** (2007), no. 2, 560 – 577, Computational Algebra.

[SX10] Bernd Sturmfels and Zhiqiang Xu, *Sagbi bases of Cox-Nagata rings*, Journal of the European Mathematical Society **12** (2010), 429–459.

Mateusz Michałek
`wajcha2@poczta.onet.pl`
Mathematical Institute of the Polish Academy of Sciences, Św. Tomasza 30, 31-027 Kraków, Poland
Max Planck Institute for Mathematics, Vivatsgasse 7, 53111 Bonn, Germany